

Patients with retinal detachment ~~are faced by a~~ fear ~~of losing the sudden and~~ significant ~~sensory function suddenly.~~ loss of vision. The purpose of our study was to understand the ~~meaning of~~ experiences of such patients, and to clarify ~~the~~ structure and special features of their experiences ~~also.~~ 11. Eleven patients with retinal detachment ~~patients-~~ took part in an ~~interview (unstructured)~~ interview and qualitative analysis of data was performed ~~by a qualitative method~~ using a phenomenologic approach. ~~Firstly,~~ First, patients with retinal detachment ~~patients-~~ assumed ~~the abnormalities in the~~ that their sight ~~will~~ would recover rapidly. ~~But abnormalities,~~ but vision became worse. Consequently, family members or friends ~~and family could persuade~~ persuaded the patients to go to hospital. Then, they were surprised and dismayed ~~when they were said to be~~ told that they ~~have~~ had to ~~enter~~ be admitted to hospital ~~and undergo~~ for emergency ~~operations~~ surgery in order to ~~not become blinded.~~ Following ~~avoid~~ blindness. After understanding that they ~~will~~ would not become blind ~~as far as receiving if they~~ underwent the operation, ~~they the patients were~~ reassured and ~~agreed~~ readily consented to ~~the operation very easily~~ surgery. However, ~~while the patients were~~ while awaiting ~~for the~~ operation ~~it started worrying,~~ patients began to worry about the possibility of going ~~blinded~~ blind and ~~felt~~ became uneasy ~~once~~ again. ~~Following~~ After the operation, ~~when~~ the patients ~~realized~~ were relieved that their ~~sights~~ sight could be maintained,

~~they relieved.~~ In order to confirm that ~~it continued to~~ progress ~~well following~~ was continuing after the operation ~~the~~, patients checked their own vision constantly.

Following ~~hospital~~ discharge from hospital, the patients ~~could return~~ returned to their previous ~~life styles~~ lifestyle, but they ~~always~~ continued to ~~worry~~ be concerned about

relapse. Eventually, the patients showed coping ~~behaviours, receiving all of~~ behavior and underwent regular ophthalmological examinations. It was found that patients with

retinal detachment ~~patients~~ experience both anxiety and relief ~~that can be changeable on,~~ which changes during the course of treatment from diagnosis to recovery.